

RESTRICTED USE PESTICIDE - Due to very high acute toxicity to Humans and Birds

For retail sale to and use only by certified applicator or persons under their direct supervision and only for those uses covered by the certified applicator's certification. Direct supervision for this product is defined as the certified applicator being physically present during application, mixing, loading, repair and cleaning of application equipment. Commercial certified applicators must also ensure that all persons involved in these activities are informed of the precautionary statements.

SECTION 24 (C) REGISTRATION FOR DISTRIBUTION AND USE ONLY IN IDAHO

METHYL PARATHION 4 EC

Emulsifiable Concentrate - An Agricultural Insecticide

EPA Reg. No. 4787 - 11-67760

EPA SLN No. ID-940005

ALL APPLICABLE DIRECTIONS, RESTRICTIONS, AND PRECAUTIONS ON THE EPA REGISTERED LABEL ARE TO BE FOLLOWED. THIS LABEL MUST BE IN THE POSSESSION OF THE USER AT THE TIME OF APPLICATION.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in any manner inconsistent with its labeling.

This label restricts use to Canola / Rapeseed oil seed varieties only.

CANOLA / RAPESEED: For control of **Cabbage Seedpod Weevil** (*Ceutorhynchus assimilis*), **Flea Beetle** (*Phyllotreta* sp.), **Diamondback Moth** (*Plutella xylostella*) and **Cabbage Aphids** (*Brevicoryne* sp.), apply 1 pint METHYL PARATHION 4E per acre by ground or air in sufficient water for coverage (minimum of 3 gallons per acre by air). Do not apply more than two applications per season. Make first application when two or more insects may be found per plant on young pods. Methyl Parathion should be applied when 10% of the bloom is remaining, or when most of the early developed pods are 1.00 - 1.25 inches in length. For **Flea Beetle** treat in early season when significant evidence of shotholing by the adults becomes evident on the seedlings.

For **Diamondback Moth larvae**, treat about mid-growing season when populations reach damaging levels and there is evidence of appreciable feeding beginning on the flowers and / or pods.

Do not apply this product through any type of irrigation system.

To protect pollinators, apply only in the evening or early morning (this includes a period from 3 hours before sunset until three hours after sunrise the following day). Do not apply within 28 days of harvest. Do not graze treated fields or feed treated forage or threshings to livestock.

Do not enter treated fields for 48 hours after application.

Cheminova Agro 1995

Manufactured For:

CHEMINOVA, INC.
1700 Route 23, Suite 210
Wayne, NJ 07470
Tel: (201) 305-6600
FAX: (201) 305-1382